Hybrid All-Pay and Winner-Pay Contests Seminar at DICE in Düsseldorf, June 5, 2018

Johan N. M. Lagerlöf Dept. of Economics, U. of Copenhagen Email: johan.lagerlof@econ.ku.dk Website: www.johanlagerlof.com

June 2, 2018

Introduction: What is a hybrid contest? (1/2)

A hybrid contest:

- In some economic, social, or political situation, each one of a number of economic agents try to win an indivisible prize.
- To increase her probability of winning, each contestant makes both **all-pay investments** and **winner-pay investments**.

Example: The competitive bidding to host the Olympic games.

- All-pay investments: Candidate cities spend money upfront, with the goal of persuading members of the IOC.
- Winner-pay investments: A city commits to build new stadia and invest in safety arrangements if being awarded the Games.
- To fix ideas, consider the following formalization:
 - Contestant *i* chooses $x_i \ge 0$ and $y_i \ge 0$ to maximize

$$\pi_i = (v_i - y_i) p_i (s_1, s_2, \dots s_n) - x_i,$$

subject to $s_i = f(x_i, y_i)$.

イロト イポト イヨト イヨ

Introduction: What is a hybrid contest? (1/2)

A hybrid contest:

- In some economic, social, or political situation, each one of a number of economic agents try to win an indivisible prize.
- To increase her probability of winning, each contestant makes both **all-pay investments** and **winner-pay investments**.

• Example: The competitive bidding to host the Olympic games.

- All-pay investments: Candidate cities spend money upfront, with the goal of persuading members of the IOC.
- Winner-pay investments: A city commits to build new stadia and invest in safety arrangements if being awarded the Games.

To fix ideas, consider the following formalization:

• Contestant *i* chooses $x_i \ge 0$ and $y_i \ge 0$ to maximize

$$\pi_i = (v_i - y_i) p_i (s_1, s_2, \dots s_n) - x_i,$$

subject to $s_i = f(x_i, y_i)$.

イロト イポト イヨト イヨ

Introduction: What is a hybrid contest? (1/2)

A hybrid contest:

- In some economic, social, or political situation, each one of a number of economic agents try to win an indivisible prize.
- To increase her probability of winning, each contestant makes both **all-pay investments** and **winner-pay investments**.

• Example: The competitive bidding to host the Olympic games.

- All-pay investments: Candidate cities spend money upfront, with the goal of persuading members of the IOC.
- Winner-pay investments: A city commits to build new stadia and invest in safety arrangements if being awarded the Games.
- To fix ideas, consider the following formalization:
 - Contestant *i* chooses $x_i \ge 0$ and $y_i \ge 0$ to maximize

$$\pi_i = (v_i - y_i) p_i (s_1, s_2, \dots s_n) - x_i,$$

subject to $s_i = f(x_i, y_i)$.

Introduction: Other examples (2/2)

- Competition for a government contract or grant:
 - *All-pay investments*: Time/effort spent on preparing proposal.
 - *Winner-pay investments*: Commit to ambitious customer service.
- A political election:
 - All-pay investments: Campaign expenditures.
 - Winner-pay investments: Electoral promises (costly if they deviate from the politician's own ideal policy).
- Rent seeking to win monopoly rights of a regulated market:
 - *All-pay investments*: Ex ante bribes (how Tullock modeled it).
 - Winner-pay investments: Conditional bribes.
- Tullock's motivation:
 - Empirical studies in the 1950s: DWL appears to be tiny.
 - Tullock: Maybe a part of profits adds to the cost of monopoly.

Literature Review (1/2)

Two earlier papers that model a hybrid contest:

- Haan and Schonbeek (2003).
 - They assume Cobb-Douglas—which here is quite restrictive.

Melkoyan (2013).

- **CES** but with $\sigma \ge 1$. Symmetric model. Hard to check SOC.
- My analysis: (i) other approach which yields easy-to-check existence condition; (ii) assumes general production function and CSF; (iii) studies both symmetric and asymmetric models.

Other contest models with more than one influence channel:

- **Sabotage in contests** (improve own performance and sabotage the others performance): Konrad (2000), Chen (2003).
- War and conflict (choice of production and appropriation): Hirschleifer (1991) and Skaperdas and Syroploulos (1997).
- Multiple all-pay "arms" (maybe with different costs): Arbatskaya and Mialon (2010).

Literature Review (1/2)

Two earlier papers that model a hybrid contest:

- Haan and Schonbeek (2003).
 - They assume Cobb-Douglas—which here is quite restrictive.

Melkoyan (2013).

- **EXAMPLE** CES but with $\sigma \geq 1$. Symmetric model. Hard to check SOC.
- My analysis: (i) other approach which yields easy-to-check existence condition; (ii) assumes general production function and CSF; (iii) studies both symmetric and asymmetric models.

• Other contest models with more than one influence channel:

- **Sabotage in contests** (improve own performance and sabotage the others performance): Konrad (2000), Chen (2003).
- War and conflict (choice of production and appropriation): Hirschleifer (1991) and Skaperdas and Syroploulos (1997).
- Multiple all-pay "arms" (maybe with different costs): Arbatskaya and Mialon (2010).

Literature Review (2/2)

Multidimensional (procurement) auctions:

• Che (2003), Branck (1997), Asker and Cantillon (2008).

- Firms bid on both price and (many dimensions of) quality.
- The components of each bid jointly determine a score.
- Auctioneer chooses bidder with highest score.

Differences:

- In their models, not both all-pay and winner-pay ingredients.
- Not a probabilistic CSF.

Optimal design of a research contest: Che and Gale (2003).

- A principal wants to procure an innovation.
- Fimrs choose both quality of innovation and the prize if winning.
- Thus, effectively, both all-pay and winner-pay ingredients
- Differences: Not a probabilistic CSF (so mixed strategy eq.), linear production function, mechanism design approach.

Literature Review (2/2)

Multidimensional (procurement) auctions:

• Che (2003), Branck (1997), Asker and Cantillon (2008).

- Firms bid on both price and (many dimensions of) quality.
- The components of each bid jointly determine a score.
- Auctioneer chooses bidder with highest score.

Differences:

- In their models, not both all-pay and winner-pay ingredients.
- Not a probabilistic CSF.

• Optimal design of a research contest: Che and Gale (2003).

- A principal wants to procure an innovation.
- Fimrs choose both quality of innovation and the prize if winning.
- Thus, effectively, both all-pay and winner-pay ingredients.
- Differences: Not a probabilistic CSF (so mixed strategy eq.), linear production function, mechanism design approach.

A model of a hybrid contest (1/2)

- $n \ge 2$ contestants try to win an indivisible prize.
- Contestant *i* chooses x_i ≥ 0 and y_i ≥ 0 to maximize the following expected payoff:

$$\pi_i = (v_i - y_i) p_i(\mathbf{s}) - x_i, \qquad \text{subject to } s_i = f(x_i, y_i),$$

where $\mathbf{s} = (s_1, s_2, \dots, s_n)$ and $s_i \ge 0$ is contestant *i*'s *score*.

- $v_i > 0$ is *i*'s valuation of the prize.
- **p**_{*i*}(**s**) is *i*'s prob. of winning (or contest success function, CSF).
- x_i is the **all-pay investment**: paid whether i wins or not.
- *y_i* is the **winner-pay investment**: paid i.f.f. *i* wins.
- It is a one-shot game where the contestants choose their investments (x_i, y_i) simultaneously with each other.

・ロト ・ 母 ト ・ ヨ ト ・ ヨ ト … ヨ

A model of a hybrid contest (2/2)

- Assumptions about $p_i(\mathbf{s})$:
 - Twice continuously differentiable in its arguments.
 - Strictly increasing and strictly concave in s_i .
 - Strictly decreasing in s_j for all $j \neq i$.
 - The contest is won by someone: $\sum_{j=1}^{n} p_j(\mathbf{s}) = 1$.
 - Later I assume that $p_i(\mathbf{s})$ is homogeneous in \mathbf{s} .

• Assumptions about $f(x_i, y_i)$:

- Thrice continuously differentiable in its arguments.
- Strictly increasing in each of its arguments.
- Strictly quasiconcave.
- Homogeneous of degree t > 0: $\forall k > 0$ $f(kx_i, ky_i) = k^t f(x_i, y_i)$.
- Inada conditions to rule out $x_i = 0$ or $y_i = 0$.

Examples:

$$p_i(\mathbf{s}) = \frac{W_i s_i^r}{\sum_{j=1}^n W_j s_j^r}, \qquad f(x_i, y_i) = \left[\alpha x^{\frac{\sigma-1}{\sigma}} + (1-\alpha) y^{\frac{\sigma-1}{\sigma}}\right]^{\frac{t\sigma}{\sigma-1}}$$

A model of a hybrid contest (2/2)

- Assumptions about $p_i(\mathbf{s})$:
 - Twice continuously differentiable in its arguments.
 - Strictly increasing and strictly concave in s_i .
 - Strictly decreasing in s_j for all $j \neq i$.
 - The contest is won by someone: $\sum_{j=1}^{n} p_j(\mathbf{s}) = 1$.
 - Later I assume that $p_i(\mathbf{s})$ is homogeneous in \mathbf{s} .
- Assumptions about $f(x_i, y_i)$:
 - Thrice continuously differentiable in its arguments.
 - Strictly increasing in each of its arguments.
 - Strictly quasiconcave.
 - Homogeneous of degree t > 0: $\forall k > 0$ $f(kx_i, ky_i) = k^t f(x_i, y_i)$.
 - Inada conditions to rule out $x_i = 0$ or $y_i = 0$.

Examples:

$$p_i(\mathbf{s}) = \frac{W_i S_i^r}{\sum_{j=1}^n W_j S_j^r}, \qquad f(x_i, y_i) = \left[\alpha x^{\frac{\sigma-1}{\sigma}} + (1-\alpha) y^{\frac{\sigma-1}{\sigma}}\right]^{\frac{t\sigma}{\sigma-1}}$$

A model of a hybrid contest (2/2)

- Assumptions about $p_i(\mathbf{s})$:
 - Twice continuously differentiable in its arguments.
 - Strictly increasing and strictly concave in s_i .
 - Strictly decreasing in s_j for all $j \neq i$.
 - The contest is won by someone: $\sum_{j=1}^{n} p_j(\mathbf{s}) = 1$.
 - Later I assume that $p_i(\mathbf{s})$ is homogeneous in \mathbf{s} .
- Assumptions about $f(x_i, y_i)$:
 - Thrice continuously differentiable in its arguments.
 - Strictly increasing in each of its arguments.
 - Strictly quasiconcave.
 - Homogeneous of degree t > 0: $\forall k > 0$ $f(kx_i, ky_i) = k^t f(x_i, y_i)$.
 - Inada conditions to rule out $x_i = 0$ or $y_i = 0$.
- Examples:

$$p_i(\mathbf{s}) = \frac{w_i s_i^r}{\sum_{j=1}^n w_j s_j^r}, \qquad f(x_i, y_i) = \left[\alpha x^{\frac{\sigma-1}{\sigma}} + (1-\alpha) y^{\frac{\sigma-1}{\sigma}}\right]^{\frac{t\sigma}{\sigma-1}}$$

Analysis (1/7)

• One possible approach:

- Plug the production function into the CSF.
- Take FOCs w.r.t. x_i and y_i .
- Used by Haan and Schoonbeek (2003) and Melkoyan (2013), assuming Cobb-Douglas and CES, respectively.

My approach: Solve for contestant *i*'s best reply in two steps:
 Compute the conditional factor demands

That is, derive optimal x_i and y_i , given **s** (so also given s_i).

- Plug the factor demands into the payoff and then characterize contestant *i*'s optimal score s_i (given s_{-i}).
- Important advantage: a single choice variable at 2, so easier to determine what conditions are required for equilibrium existence.

ヘロト ヘヨト ヘヨト

Analysis (1/7)

• One possible approach:

- Plug the production function into the CSF.
- **Take FOCs w.r.t.** x_i and y_i .
- Used by Haan and Schoonbeek (2003) and Melkoyan (2013), assuming Cobb-Douglas and CES, respectively.
- My approach: Solve for contestant *i*'s best reply in two steps:
 - **1** Compute the conditional factor demands.
 - **That is, derive optimal** x_i and y_i , given **s** (so also given s_i).
 - Plug the factor demands into the payoff and then characterize contestant *i*'s optimal score s_i (given s_{-i}).
- Important advantage: a single choice variable at 2, so easier to determine what conditions are required for equilibrium existence.

ヘロト ヘ通ト ヘヨト ヘヨト

- Contestant *i* solves (for fixed p_i): min_{$x_i,y_i} <math>p_i y_i + x_i$, subject to $f(x_i, y_i) = s_i$.</sub>
- The first-order conditions (λ_i is the Lagrange multiplier):

$$\frac{\partial \mathcal{L}_i}{\partial x_i} = 1 - \lambda_i f_1(x_i, y_i) = 0, \qquad \frac{\partial \mathcal{L}_i}{\partial y_i} = p_i - \lambda_i f_2(x_i, y_i) = 0.$$

$$\frac{1}{p_i} = \frac{f_1(x_i, y_i)}{f_2(x_i, y_i)} \stackrel{\text{def}}{=} g\left(\frac{x_i}{y_i}\right) \Rightarrow x_i = y_i h\left(\frac{1}{p_i}\right),$$

where *h* is the inverse of *g* (i.e., $h \stackrel{\text{def}}{=} g^{-1}$).

By plugging back into $s_i = f(x_i, y_i)$ and rewriting, we obtain:

$$Y_{i}(s_{i},p_{i})=\left[\frac{s_{i}}{f\left(h\left(1/p_{i}\right),1\right)}\right]^{\frac{1}{t}}, \quad X_{i}(s_{i},p_{i})=Y_{i}(s_{i},p_{i})h\left(\frac{1}{p_{i}}\right).$$

• Contestant *i*'s payoff: $\pi_i(\mathbf{s}) = p_i(\mathbf{s}) v_i - C_i[s_i, p_i(\mathbf{s})]$, where $C_i[s_i, p_i(\mathbf{s})] \stackrel{\text{def}}{=} p_i(\mathbf{s}) Y_i[s_i, p_i(\mathbf{s})] + X_i[s_i, p_i(\mathbf{s})]$

• A Nash equilibrium of the hybrid contest:

- Contestant *i* solves (for fixed p_i): min_{$x_i,y_i} <math>p_i y_i + x_i$, subject to $f(x_i, y_i) = s_i$.</sub>
- The first-order conditions (λ_i is the Lagrange multiplier):

$$\frac{\partial \mathcal{L}_i}{\partial x_i} = 1 - \lambda_i f_1(x_i, y_i) = 0, \qquad \frac{\partial \mathcal{L}_i}{\partial y_i} = p_i - \lambda_i f_2(x_i, y_i) = 0.$$

$$\frac{1}{p_i} = \frac{f_1(x_i, y_i)}{f_2(x_i, y_i)} \stackrel{\text{def}}{=} g\left(\frac{x_i}{y_i}\right) \Rightarrow x_i = y_i h\left(\frac{1}{p_i}\right)$$

where *h* is the inverse of *g* (i.e., $h \stackrel{\text{def}}{=} g^{-1}$).

By plugging back into $s_i = f(x_i, y_i)$ and rewriting, we obtain:

$$Y_{i}\left(s_{i}, p_{i}\right) = \left[\frac{s_{i}}{f\left(h\left(1/p_{i}\right), 1\right)}\right]^{\frac{1}{t}}, \quad X_{i}\left(s_{i}, p_{i}\right) = Y_{i}\left(s_{i}, p_{i}\right)h\left(\frac{1}{p_{i}}\right).$$

Contestant *i*'s payoff: $\pi_i(\mathbf{s}) = p_i(\mathbf{s}) v_i - C_i[s_i, p_i(\mathbf{s})]$, where $C_i[s_i, p_i(\mathbf{s})] \stackrel{\text{def}}{=} p_i(\mathbf{s}) Y_i[s_i, p_i(\mathbf{s})] + X_i[s_i, p_i(\mathbf{s})]$

• A Nash equilibrium of the hybrid contest:

- Contestant *i* solves (for fixed p_i): min_{$x_i,y_i} <math>p_i y_i + x_i$, subject to $f(x_i, y_i) = s_i$.</sub>
- The first-order conditions (λ_i is the Lagrange multiplier):

$$\frac{\partial \mathcal{L}_i}{\partial x_i} = 1 - \lambda_i f_1(x_i, y_i) = 0, \qquad \frac{\partial \mathcal{L}_i}{\partial y_i} = p_i - \lambda_i f_2(x_i, y_i) = 0.$$

$$\frac{1}{p_i} = \frac{f_1(x_i, y_i)}{f_2(x_i, y_i)} \stackrel{\text{def}}{=} g\left(\frac{x_i}{y_i}\right) \Rightarrow x_i = y_i h\left(\frac{1}{p_i}\right).$$

where *h* is the inverse of *g* (i.e., $h \stackrel{\text{def}}{=} g^{-1}$).

By plugging back into $s_i = f(x_i, y_i)$ and rewriting, we obtain:

$$Y_{i}\left(s_{i}, p_{i}\right) = \left[\frac{s_{i}}{f\left(h\left(1/p_{i}\right), 1\right)}\right]^{\frac{1}{t}}, \quad X_{i}\left(s_{i}, p_{i}\right) = Y_{i}\left(s_{i}, p_{i}\right)h\left(\frac{1}{p_{i}}\right).$$

Contestant *i*'s payoff: $\pi_i(\mathbf{s}) = p_i(\mathbf{s}) V_i - C_i[s_i, p_i(\mathbf{s})]$, where

• A Nash equilibrium of the hybrid contest:

- Contestant *i* solves (for fixed p_i): min_{$x_i,y_i} <math>p_i y_i + x_i$, subject to $f(x_i, y_i) = s_i$.</sub>
- The first-order conditions (λ_i is the Lagrange multiplier):

$$\frac{\partial \mathcal{L}_i}{\partial x_i} = 1 - \lambda_i f_1(x_i, y_i) = 0, \qquad \frac{\partial \mathcal{L}_i}{\partial y_i} = p_i - \lambda_i f_2(x_i, y_i) = 0.$$

$$\frac{1}{p_i} = \frac{f_1(x_i, y_i)}{f_2(x_i, y_i)} \stackrel{\text{def}}{=} g\left(\frac{x_i}{y_i}\right) \Rightarrow x_i = y_i h\left(\frac{1}{p_i}\right)$$

where *h* is the inverse of *g* (i.e., $h \stackrel{\text{def}}{=} g^{-1}$)

By plugging back into $s_i = f(x_i, y_i)$ and rewriting, we obtain:

$$Y_{i}\left(s_{i}, p_{i}\right) = \left[\frac{s_{i}}{f\left(h\left(1/p_{i}\right), 1\right)}\right]^{\frac{1}{t}}, \quad X_{i}\left(s_{i}, p_{i}\right) = Y_{i}\left(s_{i}, p_{i}\right)h\left(\frac{1}{p_{i}}\right).$$

Contestant *i*'s payoff: $\pi_i(\mathbf{s}) = p_i(\mathbf{s}) v_i - C_i[s_i, p_i(\mathbf{s})]$, where

A Nash equilibrium of the hybrid contest:

- Contestant *i* solves (for fixed p_i): $\min_{x_i, y_i} p_i y_i + x_i$, subject to $f(x_i, y_i) = s_i$.
- The first-order conditions (λ_i is the Lagrange multiplier):

$$\frac{\partial \mathcal{L}_i}{\partial x_i} = 1 - \lambda_i f_1(x_i, y_i) = 0, \qquad \frac{\partial \mathcal{L}_i}{\partial y_i} = p_i - \lambda_i f_2(x_i, y_i) = 0.$$

$$\frac{1}{p_i} = \frac{f_1(x_i, y_i)}{f_2(x_i, y_i)} \stackrel{\text{def}}{=} g\left(\frac{x_i}{y_i}\right) \Rightarrow x_i = y_i h\left(\frac{1}{p_i}\right),$$

where *h* is the inverse of *g* (i.e., $h \stackrel{\text{def}}{=} g^{-1}$).

By plugging back into $s_i = f(x_i, y_i)$ and rewriting, we obtain:

$$Y_{i}\left(s_{i}, p_{i}\right) = \left[\frac{s_{i}}{f\left(h\left(1/p_{i}\right), 1\right)}\right]^{\frac{1}{t}}, \quad X_{i}\left(s_{i}, p_{i}\right) = Y_{i}\left(s_{i}, p_{i}\right)h\left(\frac{1}{p_{i}}\right).$$

Contestant *i*'s payoff: $\pi_i(\mathbf{s}) = p_i(\mathbf{s}) v_i - C_i[s_i, p_i(\mathbf{s})]$, where

 $C_{i}[s_{i}, p_{i}(\mathbf{s})] \stackrel{\text{def}}{=} p_{i}(\mathbf{s}) Y_{i}[s_{i}, p_{i}(\mathbf{s})] + X_{i}[s_{i}, p_{i}(\mathbf{s})]$

• A Nash equilibrium of the hybrid contest:

- Contestant *i* solves (for fixed p_i): min_{$x_i,y_i} <math>p_i y_i + x_i$, subject to $f(x_i, y_i) = s_i$.</sub>
- The first-order conditions (λ_i is the Lagrange multiplier):

$$\frac{\partial \mathcal{L}_i}{\partial x_i} = 1 - \lambda_i f_1(x_i, y_i) = 0, \qquad \frac{\partial \mathcal{L}_i}{\partial y_i} = p_i - \lambda_i f_2(x_i, y_i) = 0.$$

$$\frac{1}{p_i} = \frac{f_1(x_i, y_i)}{f_2(x_i, y_i)} \stackrel{\text{def}}{=} g\left(\frac{x_i}{y_i}\right) \Rightarrow x_i = y_i h\left(\frac{1}{p_i}\right),$$

where *h* is the inverse of *g* (i.e., $h \stackrel{\text{def}}{=} g^{-1}$).

By plugging back into $s_i = f(x_i, y_i)$ and rewriting, we obtain:

$$Y_{i}(s_{i},p_{i}) = \left[\frac{s_{i}}{f(h(1/p_{i}),1)}\right]^{\frac{1}{t}}, \quad X_{i}(s_{i},p_{i}) = Y_{i}(s_{i},p_{i})h\left(\frac{1}{p_{i}}\right)$$


• Contestant *i*'s payoff: $\pi_i(\mathbf{s}) = p_i(\mathbf{s}) v_i - C_i[s_i, p_i(\mathbf{s})]$, where $C_i[s_i, p_i(\mathbf{s})] \stackrel{\text{def}}{=} p_i(\mathbf{s}) Y_i[s_i, p_i(\mathbf{s})] + X_i[s_i, p_i(\mathbf{s})]$.

A Nash equilibrium of the hybrid contest:

• A profile **s**^{*} such that $\pi_i(\mathbf{s}^*) \ge \pi_i(s_i, \mathbf{s}^*_{-\mathbf{i}})$, all i and all $s_i \ge 0$

Analysis (3/7)

The cost-minimization problem and the *h* function



Define the following elasticities:

• The elasticity of output w.r.t. x_i : $\eta\left(\frac{1}{p_i}\right) \stackrel{\text{def}}{=} \frac{f_1\left[h\left(\frac{1}{p_i}\right), 1\right]h\left(\frac{1}{p_i}\right)}{f\left[h\left(\frac{1}{p_i}\right), 1\right]}$.

• The elasticity of substitution: $\sigma\left(\frac{1}{p_i}\right) \stackrel{\text{\tiny def}}{=} -\frac{h'\left(\frac{1}{p_i}\right)\frac{1}{p_i}}{h\left(\frac{1}{p_i}\right)}$.

The elasticity of the win probability w.r.t. s_i: ε_i (s) ^{def} ∂p_i s_i/∂s_i p_i.
 We have that η ∈ (0, t), σ > 0, and ε_i ∈ (0, 1).

Assumption 1. The production function and the CSF satisfy: (i) $t \le 1$ and $\varepsilon_i(\mathbf{s}) \eta\left(\frac{1}{p_i}\right) \sigma\left(\frac{1}{p_i}\right) \le 2$ (for all p_i and \mathbf{s});

Proposition 1. Suppose Assumption 1 is satisfied. Then there
exists a pure strategy Nash equilibrium of the hybrid contest.

Define the following elasticities:

• The elasticity of output w.r.t. x_i : $\eta\left(\frac{1}{p_i}\right) \stackrel{\text{def}}{=} \frac{f_1\left[h\left(\frac{1}{p_i}\right), 1\right]h\left(\frac{1}{p_i}\right)}{f\left[h\left(\frac{1}{p_i}\right), 1\right]}$.

• The elasticity of substitution: $\sigma\left(\frac{1}{p_i}\right) \stackrel{\text{def}}{=} -\frac{h'\left(\frac{1}{p_i}\right)\frac{1}{p_i}}{h\left(\frac{1}{p_i}\right)}$.

The elasticity of the win probability w.r.t. s_i: ε_i (s) ^{def} ∂ p_i s_i / ∂s_i ρ_i.
 We have that η ∈ (0, t), σ > 0, and ε_i ∈ (0, 1).

Assumption 1. The production function and the CSF satisfy: (i) $t \le 1$ and $\varepsilon_i(\mathbf{s}) \eta\left(\frac{1}{p_i}\right) \sigma\left(\frac{1}{p_i}\right) \le 2$ (for all p_i and \mathbf{s});

Proposition 1. Suppose Assumption 1 is satisfied. Then there
exists a pure strategy Nash equilibrium of the hybrid contest.

Define the following elasticities:

The elasticity of output w.r.t. x_i: η (1/p_i) = f_1[h(1/p_i),1]h(1/p_i))/f[h(1/p_i),1].
 The elasticity of substitution: σ (1/p_i) = -h'(1/p_i)/1/h(1/p_i).

The elasticity of the win probability w.r.t. s_i: ε_i (s) ^{def} ∂p_i s_i / ∂s_i p_i.
 We have that η ∈ (0, t), σ > 0, and ε_i ∈ (0, 1).

Assumption 1. The production function and the CSF satisfy:

 (i) t ≤ 1 and ε_i(s) η (1/p_i) σ (1/p_i) ≤ 2
 (for all p_i and s);

 Proposition 1. Suppose Assumption 1 is satisfied. Then there


exists a pure strategy Nash equilibrium of the hybrid contest.

Define the following elasticities:

The elasticity of output w.r.t. x_i: η (1/p_i) = f₁[h(1/p_i),1]h(1/p_i)/f[h(1/p_i),1].
 The elasticity of substitution: σ (1/p_i) = -h'(1/p_i)1/p_i/h(1/q_i).

- The elasticity of the win probability w.r.t. s_i: ε_i (s) ^{def} ∂ p_i s_i / ∂s_i ρ_i.
 We have that η ∈ (0, t), σ > 0, and ε_i ∈ (0, 1).
- Assumption 1. The production function and the CSF satisfy: (i) $t \le 1$ and $\varepsilon_i(\mathbf{s}) \eta\left(\frac{1}{p_i}\right) \sigma\left(\frac{1}{p_i}\right) \le 2$ (for all p_i and \mathbf{s});
- Proposition 1. Suppose Assumption 1 is satisfied. Then there exists a pure strategy Nash equilibrium of the hybrid contest.

Assume a CES production function, t = 1, $r \le 1$, and



To check the SOC with Melkoyan's analytical approach is cumbersome and in the end he relies on numerical simulations:

[...] one can demonstrate, after a series of tedious algebraic manipulations, that a player's payoff function is locally concave at the symmetric equilibrium candidate in (7) if and only if [large mathematical expression]. [...] Numerical simulations indicate that this inequality is violated only for extreme values of the parameters [...]. In addition to verifying the local second-order conditions, I have used numerical simulations to verify that the global second-order conditions are satisfied under a wide range of scenarios.

イロト イポト イヨト イヨト 二日

Recall:
$$\pi_i(\mathbf{s}) = p_i(\mathbf{s}) v_i - C_i[s_i, p_i(\mathbf{s})].$$

• The FOC (with an equality if $s_i > 0$):

$$\frac{\partial \pi_{i}\left(\mathbf{s}\right)}{\partial s_{i}}=\frac{\partial p_{i}\left(\mathbf{s}\right)}{\partial s_{i}}v_{i}-C_{1}\left(s_{i},p_{i}\right)-C_{2}\left(s_{i},p_{i}\right)\frac{\partial p_{i}\left(\mathbf{s}\right)}{\partial s_{i}}\leq0.$$

• Use Shephard's lemma, $C_2(s_i, p_i) = Y_i[s_i, p_i(\mathbf{s})]$:

$$\left[v_{i}-Y_{i}\left(s_{i},p_{i}\left(\mathbf{s}\right)\right)\right]\frac{\partial p_{i}\left(\mathbf{s}\right)}{\partial s_{i}}\leq C_{1}\left(s_{i},p_{i}\right),$$
(1)

June 5, 2018

14 / 24

with an equality if $s_i > 0$.

Proposition 2. Suppose Assumption 1 is satisfied. Then
 s^{*} = (s₁^{*},..., s_n^{*}) is a pure strategy Nash equilibrium of the hybrid contest if and only if condition (1) holds, with equality if s_i^{*} > 0, for each contestant *i*.

- **Recall:** $\pi_i(\mathbf{s}) = p_i(\mathbf{s}) v_i C_i[s_i, p_i(\mathbf{s})].$
- The FOC (with an equality if $s_i > 0$):

$$\frac{\partial \pi_{i}\left(\mathbf{s}\right)}{\partial s_{i}}=\frac{\partial p_{i}\left(\mathbf{s}\right)}{\partial s_{i}}v_{i}-C_{1}\left(s_{i},p_{i}\right)-C_{2}\left(s_{i},p_{i}\right)\frac{\partial p_{i}\left(\mathbf{s}\right)}{\partial s_{i}}\leq0.$$

• Use Shephard's lemma, $C_2(s_i, p_i) = Y_i[s_i, p_i(\mathbf{s})]$:

$$\left[v_{i}-Y_{i}\left(s_{i},p_{i}\left(\mathbf{s}\right)\right)\right]\frac{\partial p_{i}\left(\mathbf{s}\right)}{\partial s_{i}}\leq C_{1}\left(s_{i},p_{i}\right),\tag{1}$$

with an equality if $s_i > 0$.

Proposition 2. Suppose Assumption 1 is satisfied. Then
 s* = (s₁*,..., s_n*) is a pure strategy Nash equilibrium of the hybrid contest if and only if condition (1) holds, with equality if s_i* > 0, for each contestant *i*.

- **Recall:** $\pi_i(\mathbf{s}) = p_i(\mathbf{s}) v_i C_i[s_i, p_i(\mathbf{s})].$
- The FOC (with an equality if $s_i > 0$):

$$\frac{\partial \pi_{i}\left(\mathbf{s}\right)}{\partial s_{i}}=\frac{\partial p_{i}\left(\mathbf{s}\right)}{\partial s_{i}}v_{i}-C_{1}\left(s_{i},p_{i}\right)-C_{2}\left(s_{i},p_{i}\right)\frac{\partial p_{i}\left(\mathbf{s}\right)}{\partial s_{i}}\leq0.$$

• Use Shephard's lemma, $C_2(s_i, p_i) = Y_i[s_i, p_i(\mathbf{s})]$:

$$\left[v_{i}-Y_{i}\left(s_{i},p_{i}\left(\mathbf{s}\right)\right)\right]\frac{\partial p_{i}\left(\mathbf{s}\right)}{\partial s_{i}}\leq C_{1}\left(s_{i},p_{i}\right),$$
(1)

with an equality if $s_i > 0$.

Proposition 2. Suppose Assumption 1 is satisfied. Then
 s* = (s₁*,..., s_n*) is a pure strategy Nash equilibrium of the hybrid contest if and only if condition (1) holds, with equality if s_i* > 0, for each contestant *i*.

Recall:
$$\pi_i(\mathbf{s}) = p_i(\mathbf{s}) v_i - C_i[s_i, p_i(\mathbf{s})].$$

• The FOC (with an equality if $s_i > 0$):

$$\frac{\partial \pi_{i}\left(\mathbf{s}\right)}{\partial s_{i}}=\frac{\partial p_{i}\left(\mathbf{s}\right)}{\partial s_{i}}v_{i}-C_{1}\left(s_{i},p_{i}\right)-C_{2}\left(s_{i},p_{i}\right)\frac{\partial p_{i}\left(\mathbf{s}\right)}{\partial s_{i}}\leq0.$$

Use Shephard's lemma, $C_2(s_i, p_i) = Y_i[s_i, p_i(\mathbf{s})]$:

$$\left[\mathbf{v}_{i}-Y_{i}\left(s_{i},p_{i}\left(\mathbf{s}\right)\right)\right]\frac{\partial p_{i}\left(\mathbf{s}\right)}{\partial s_{i}}\leq C_{1}\left(s_{i},p_{i}\right), \tag{1}$$

with an equality if $s_i > 0$.

Proposition 2. Suppose Assumption 1 is satisfied. Then
 s* = (s₁*,...,s_n*) is a pure strategy Nash equilibrium of the hybrid contest if and only if condition (1) holds, with equality if s_i* > 0, for each contestant *i*.

• Recall:
$$\pi_i(\mathbf{s}) = p_i(\mathbf{s}) v_i - C_i[s_i, p_i(\mathbf{s})].$$

• The FOC (with an equality if $s_i > 0$):

$$\frac{\partial \pi_{i}\left(\mathbf{s}\right)}{\partial s_{i}}=\frac{\partial p_{i}\left(\mathbf{s}\right)}{\partial s_{i}}v_{i}-C_{1}\left(s_{i},p_{i}\right)-C_{2}\left(s_{i},p_{i}\right)\frac{\partial p_{i}\left(\mathbf{s}\right)}{\partial s_{i}}\leq0.$$

• Use Shephard's lemma, $C_2(s_i, p_i) = Y_i[s_i, p_i(\mathbf{s})]$:

$$\left[\mathbf{v}_{i}-Y_{i}\left(\mathbf{s}_{i},\mathbf{p}_{i}\left(\mathbf{s}\right)\right)\right]\frac{\partial \mathbf{p}_{i}\left(\mathbf{s}\right)}{\partial \mathbf{s}_{i}}\leq C_{1}\left(\mathbf{s}_{i},\mathbf{p}_{i}\right), \tag{1}$$

with an equality if $s_i > 0$.

Proposition 2. Suppose Assumption 1 is satisfied. Then
 s^{*} = (s^{*}₁,..., s^{*}_n) is a pure strategy Nash equilibrium of the hybrid contest if and only if condition (1) holds, with equality if s^{*}_i > 0, for each contestant *i*.

A Symmetric Hybrid Contest (1/4)

Assumption 2. The CSF is symmetric and homogeneous of degree 0.Note that, thanks to Assumption 2:

 $\frac{\partial p_i(s, s, \dots, s)}{\partial s_i} = \frac{\widehat{\varepsilon}(n)}{ns}, \text{ where } \widehat{\varepsilon}(n) \stackrel{\text{def}}{=} \varepsilon_i (1, 1, \dots, 1).$ $\blacksquare \text{ Use this in the FOC and impose symmetry:}$ $(v - y^*) \frac{\widehat{\varepsilon}(n)}{ns^*} = C_1 \left[s^*, \frac{1}{n} \right] = \frac{1}{ts^*} C \left[s^*, \frac{1}{n} \right] = \frac{1}{ts^*} \left[\frac{y^*}{n} + x^* \right]$ $\Leftrightarrow (v - y^*) t\widehat{\varepsilon}(n) = y^* + nx^*. \text{ And from before, } x^* = h(n)y^*.$ $\blacksquare \text{ The last equalities are linear in } x^* \text{ and } y^*, \text{ so easy to solve.}$ $\blacksquare \text{$ **Proposition 3.**Within the family of sym. eq., there is a unique set of the set of

pure strategy equilibrium: $s^* = f[h(n),1]\left(y^*
ight)^t$, $x^* = h(n)y^*$, and

$$y^* = \frac{t\widehat{\varepsilon}(n)v}{1 + nh(n) + t\widehat{\varepsilon}(n)}.$$

A Symmetric Hybrid Contest (1/4)

Assumption 2. The CSF is symmetric and homogeneous of degree 0.Note that, thanks to Assumption 2:

$$\frac{\partial p_i(s, s, \dots, s)}{\partial s_i} = \frac{\widehat{\varepsilon}(n)}{ns}, \text{ where } \widehat{\varepsilon}(n) \stackrel{\text{def}}{=} \varepsilon_i (1, 1, \dots, 1).$$
Use this in the FOC and impose symmetry:

$$(v - y^*)\frac{\widehat{\varepsilon}(n)}{ns^*} = C_1\left[s^*, \frac{1}{n}\right] = \frac{1}{ts^*}C\left[s^*, \frac{1}{n}\right] = \frac{1}{ts^*}\left[\frac{y^*}{n} + x^*\right]$$

 $\Leftrightarrow (v - y^*) t\widehat{\varepsilon}(n) = y^* + nx^*.$ And from before, $x^* = h(n)y^*.$ The last equalities are linear in x^* and y^* , so easy to solve.

Proposition 3. Within the family of sym. eq., there is a unique pure strategy equilibrium: $s^* = f[h(n), 1] (y^*)^t$, $x^* = h(n)y^*$, and

$$y^* = \frac{t\widehat{\varepsilon}(n)v}{1 + nh(n) + t\widehat{\varepsilon}(n)}.$$

A Symmetric Hybrid Contest (1/4)

Assumption 2. The CSF is symmetric and homogeneous of degree 0.Note that, thanks to Assumption 2:

$$\frac{\partial p_i(s, s, \dots, s)}{\partial s_i} = \frac{\widehat{\varepsilon}(n)}{ns}, \text{ where } \widehat{\varepsilon}(n) \stackrel{\text{def}}{=} \varepsilon_i (1, 1, \dots, 1).$$
Use this in the FOC and impose symmetry:

$$(v-y^*)\frac{\widehat{\varepsilon}(n)}{ns^*} = C_1\left[s^*, \frac{1}{n}\right] = \frac{1}{ts^*}C\left[s^*, \frac{1}{n}\right] = \frac{1}{ts^*}\left[\frac{y^*}{n} + x^*\right]$$

 $\Leftrightarrow (v - y^*) t\widehat{\varepsilon}(n) = y^* + nx^*. \text{ And from before, } x^* = h(n)y^*.$ The last equalities are linear in x^* and y^* , so easy to solve.

Proposition 3. Within the family of sym. eq., there is a unique pure strategy equilibrium: $s^* = f[h(n), 1] (y^*)^t$, $x^* = h(n)y^*$, and

$$y^* = \frac{t\widehat{\varepsilon}(n)v}{1 + nh(n) + t\widehat{\varepsilon}(n)}.$$

Assumption 2. The CSF is symmetric and homogeneous of degree 0.Note that, thanks to Assumption 2:

$$\frac{\partial p_i(s, s, \dots, s)}{\partial s_i} = \frac{\widehat{\varepsilon}(n)}{ns}, \text{ where } \widehat{\varepsilon}(n) \stackrel{\text{\tiny def}}{=} \varepsilon_i (1, 1, \dots, 1).$$
Use this in the FOC and impose symmetry:

$$(v-y^*)\frac{\widehat{\varepsilon}(n)}{ns^*} = C_1\left[s^*, \frac{1}{n}\right] = \frac{1}{ts^*}C\left[s^*, \frac{1}{n}\right] = \frac{1}{ts^*}\left[\frac{y^*}{n} + x^*\right]$$

 \Leftrightarrow $(v - y^*) t\hat{\varepsilon}(n) = y^* + nx^*$. And from before, $x^* = h(n)y^*$.

The last equalities are linear in x^* and y^* , so easy to solve.

$$y^* = \frac{t\widehat{\varepsilon}(n)v}{1 + nh(n) + t\widehat{\varepsilon}(n)}.$$

Assumption 2. The CSF is symmetric and homogeneous of degree 0.Note that, thanks to Assumption 2:

$$\frac{\partial p_i(s, s, \dots, s)}{\partial s_i} = \frac{\widehat{\varepsilon}(n)}{ns}, \text{ where } \widehat{\varepsilon}(n) \stackrel{\text{\tiny def}}{=} \varepsilon_i (1, 1, \dots, 1).$$
Use this in the FOC and impose symmetry:

$$(v-y^*)\frac{\widehat{\varepsilon}(n)}{ns^*} = C_1\left[s^*, \frac{1}{n}\right] = \frac{1}{ts^*}C\left[s^*, \frac{1}{n}\right] = \frac{1}{ts^*}\left[\frac{y^*}{n} + x^*\right]$$

 \Leftrightarrow $(v - y^*) t \hat{\varepsilon}(n) = y^* + nx^*$. And from before, $x^* = h(n)y^*$.

The last equalities are linear in x^* and y^* , so easy to solve.

$$y^* = \frac{t\widehat{\varepsilon}(n)v}{1 + nh(n) + t\widehat{\varepsilon}(n)}.$$

Assumption 2. The CSF is symmetric and homogeneous of degree 0.Note that, thanks to Assumption 2:

$$\frac{\partial p_i(s, s, \dots, s)}{\partial s_i} = \frac{\widehat{\varepsilon}(n)}{ns}, \text{ where } \widehat{\varepsilon}(n) \stackrel{\text{\tiny def}}{=} \varepsilon_i (1, 1, \dots, 1).$$

Use this in the FOC and impose symmetry:

$$(v-y^*)\frac{\widehat{\varepsilon}(n)}{ns^*} = C_1\left[s^*, \frac{1}{n}\right] = \frac{1}{ts^*}C\left[s^*, \frac{1}{n}\right] = \frac{1}{ts^*}\left[\frac{y^*}{n} + x^*\right]$$

 $\Leftrightarrow (\mathbf{v} - \mathbf{y}^*) t\widehat{\varepsilon}(n) = \mathbf{y}^* + n\mathbf{x}^*. \text{ And from before, } \mathbf{x}^* = h(n)\mathbf{y}^*.$

The last equalities are linear in x^* and y^* , so easy to solve.

$$y^* = \frac{t\widehat{\varepsilon}(n)v}{1 + nh(n) + t\widehat{\varepsilon}(n)}.$$

Assumption 2. The CSF is symmetric and homogeneous of degree 0.Note that, thanks to Assumption 2:

$$\frac{\partial p_i(s, s, \dots, s)}{\partial s_i} = \frac{\widehat{\varepsilon}(n)}{ns}, \text{ where } \widehat{\varepsilon}(n) \stackrel{\text{\tiny def}}{=} \varepsilon_i (1, 1, \dots, 1).$$

Use this in the FOC and impose symmetry:

$$(v-y^*)\frac{\widehat{\varepsilon}(n)}{ns^*} = C_1\left[s^*, \frac{1}{n}\right] = \frac{1}{ts^*}C\left[s^*, \frac{1}{n}\right] = \frac{1}{ts^*}\left[\frac{y^*}{n} + x^*\right]$$

 \Leftrightarrow $(v - y^*) t \widehat{\varepsilon}(n) = y^* + nx^*$. And from before, $x^* = h(n)y^*$.

The last equalities are linear in x^* and y^* , so easy to solve.

$$y^* = \frac{t\widehat{\varepsilon}(n)v}{1 + nh(n) + t\widehat{\varepsilon}(n)}.$$

Assumption 2. The CSF is symmetric and homogeneous of degree 0.Note that, thanks to Assumption 2:

$$\frac{\partial p_i(s, s, \ldots, s)}{\partial s_i} = \frac{\widehat{\varepsilon}(n)}{ns}, \text{ where } \widehat{\varepsilon}(n) \stackrel{\text{\tiny def}}{=} \varepsilon_i (1, 1, \ldots, 1).$$

Use this in the FOC and impose symmetry:

$$(v-y^*)\frac{\widehat{\varepsilon}(n)}{ns^*} = C_1\left[s^*, \frac{1}{n}\right] = \frac{1}{ts^*}C\left[s^*, \frac{1}{n}\right] = \frac{1}{ts^*}\left[\frac{y^*}{n} + x^*\right]$$

 $\Leftrightarrow (v - y^*) t\widehat{\varepsilon}(n) = y^* + nx^*.$ And from before, $x^* = h(n)y^*.$ The last equalities are linear in x^* and y^* , so easy to solve.

Proposition 3. Within the family of sym. eq., there is a unique pure strategy equilibrium: s^{*} = f[h(n), 1] (y^{*})^t, x^{*} = h(n)y^{*}, and

$$y^* = \frac{t\widehat{\varepsilon}(n)v}{1+nh(n)+t\widehat{\varepsilon}(n)}.$$

Assumption 2. The CSF is symmetric and homogeneous of degree 0.Note that, thanks to Assumption 2:

$$\frac{\partial p_i(s, s, \ldots, s)}{\partial s_i} = \frac{\widehat{\varepsilon}(n)}{ns}, \text{ where } \widehat{\varepsilon}(n) \stackrel{\text{\tiny def}}{=} \varepsilon_i (1, 1, \ldots, 1).$$

Use this in the FOC and impose symmetry:

$$(v-y^*)\frac{\widehat{\varepsilon}(n)}{ns^*} = C_1\left[s^*, \frac{1}{n}\right] = \frac{1}{ts^*}C\left[s^*, \frac{1}{n}\right] = \frac{1}{ts^*}\left[\frac{y^*}{n} + x^*\right]$$

 $\Leftrightarrow (v - y^*) t\widehat{\varepsilon}(n) = y^* + nx^*.$ And from before, $x^* = h(n)y^*.$ The last equalities are linear in x^* and y^* , so easy to solve.

$$y^* = rac{t\widehat{\varepsilon}(n)v}{1+nh(n)+t\widehat{\varepsilon}(n)}.$$

Proposition 4. Effect of more contestants on *x*^{*} and *y*^{*}:

$$rac{\partial x^*}{\partial n} < 0 \Leftrightarrow \sigma(n) > -rac{n(n-2)h(n)-1}{(n-1)[1+t\widehat{\varepsilon}(n)]}, \ rac{\partial y^*}{\partial n} > 0 \Leftrightarrow \sigma(n) > rac{n(n-2)h(n)-1}{(n-1)nh(n)};$$

and if $\sigma(n) \ge 1$, then necessarily $\frac{\partial x^*}{\partial n} < 0$ and $\frac{\partial y^*}{\partial n} > 0$. In order to understand the above:

- More contestants means a lower probability of winning.
- This lowers the relative cost of investing in y_i.
- So whenever $\sigma(n)$ is sufficiently large, $\frac{\partial y^*}{\partial n} > 0$ and $\frac{\partial x^*}{\partial n} < 0$.
- But if $\sigma(n)$ small, the derivatives must have the same sign. For:

$$\frac{\partial y^*}{\partial n}\frac{n}{y^*} = \sigma(n) + \frac{\partial x^*}{\partial n}\frac{n}{x^*} \qquad \text{(follows from } x^* = h(n)y^*\text{)}.$$

As $\sigma(n) \rightarrow 0$, the production function requires x_i and y_i to be used in fixed proportions (a Leontief production technology).

- The total amount of equilibrium expenditures in the symmetric hybrid model is defined as $R^{H} \stackrel{\text{def}}{=} nC\left[s^*, \frac{1}{n}\right]$.
- The corresponding amount in the all-pay contest: R^A = t c(n)v.
 Proposition 5, part (a). In the symmetric model:

$$R^{\mathsf{H}} = (1 - rac{y^{*}}{v})R^{\mathsf{A}} = \left[rac{1}{v\left[1 + nh(n)
ight]} + rac{1}{R^{\mathsf{A}}}
ight]^{-1}$$

In particular, for any finite *n*, we have $R^{H} < R^{A}$.

The payoff suggests the intuition: π_i = (v_i − y_i) p_i (s) − x_i.
 Proposition 5, part (b). In the symmetric model, suppose p_i(s) = φ(s_i)/∑ⁿ_{j=1} φ(s_j), where φ is a strictly increasing and concave function satisfying φ(0) = 0.

Then R^{H} is weakly increasing in *n* if and only if: (i)

$$\sigma(n) \le 1 + \frac{4n}{tr(n-1)^2}; \tag{2}$$

or (ii) inequality (2) is violated and $h(n) \notin (\Xi_L, \Xi_H)$. See figure

- The total amount of equilibrium expenditures in the symmetric hybrid model is defined as $R^{\mathsf{H}} \stackrel{\text{def}}{=} nC\left[s^*, \frac{1}{n}\right]$.
- The corresponding amount in the all-pay contest: R^A = t c(n)v.
 Proposition 5, part (a). In the symmetric model:

$$R^{\mathsf{H}} = (1 - rac{y^{*}}{v})R^{\mathsf{A}} = \left[rac{1}{v\left[1 + nh(n)
ight]} + rac{1}{R^{\mathsf{A}}}
ight]^{-1}$$

In particular, for any finite *n*, we have $R^{H} < R^{A}$.

The payoff suggests the intuition: π_i = (v_i − y_i) p_i (s) − x_i.
 Proposition 5, part (b). In the symmetric model, suppose p_i(s) = φ(s_i)/∑ⁿ_{j=1} φ(s_j), where φ is a strictly increasing and concave function satisfying φ(0) = 0.


• Then R^{H} is weakly increasing in *n* if and only if: (i)

$$\sigma(n) \le 1 + \frac{4n}{tr(n-1)^2}; \tag{2}$$

or (ii) inequality (2) is violated and $h(n) \notin (\Xi_L, \Xi_H)$. See figure!

Illustration of result (b)

• Assume CES, t = 1, and n = 10.



J. Lagerlöf (U of Copenhagen)

Hybrid All-Pay and Winner-Pay Contests

June 5, 2018 18 / 24

- I assume n = 2 and I study three models:
 - The CSF is biased in favor of one contestant.
 - One contestant has a higher valuation than the other.
 - I also endogenize the degree of bias.
- **Assumption 3.** The CSF is given by

$$p_i(\mathbf{s}) = \frac{w_i s_i^r}{w_1 s_1^r + w_2 s_2^r}.$$

The following three equations define equilibrium values of p_1^* , y_1^* , and y_2^* :

$$y_i^* = \frac{rtp_i^*(1-p_i^*)v_i}{rtp_i^*(1-p_i^*) + p_i^* + h\left(\frac{1}{p_i^*}\right)}, \quad \text{for } i = 1, 2, \text{ and } \Upsilon(p_1^*) = 0, \text{ where}$$
$$\Upsilon(p_1) \stackrel{\text{def}}{=} \frac{\frac{w_2 v_2^{rt}}{w_1 v_1^{rt}} p_1 f\left[h\left(\frac{1}{1-p_1}\right), 1\right]^r}{\left[rtp_1(1-p_1) + 1 - p_1 + h\left(\frac{1}{1-p_1}\right)\right]^{rt}} - \frac{(1-p_1) f\left[h\left(\frac{1}{p_1}\right), 1\right]^r}{\left[rtp_1(1-p_1) + p_1 + h\left(\frac{1}{p_1}\right)\right]^{rt}}.$$

The equilibrium is unique if $r\eta\left(\frac{1}{p_i}\right)\sigma\left(\frac{1}{p_i}\right) \leq 1$.

June 5, 2018 19 / 24

A B > A B > A B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A

- I assume n = 2 and I study three models:
 - The CSF is biased in favor of one contestant.
 - One contestant has a higher valuation than the other.
 - I also endogenize the degree of bias.
- **Assumption 3.** The CSF is given by

$$p_i(\mathbf{s}) = \frac{w_i s_i^r}{w_1 s_1^r + w_2 s_2^r}.$$

The following three equations define equilibrium values of p_1^* , y_1^* , and y_2^* :

$$y_{i}^{*} = \frac{rtp_{i}^{*}(1-p_{i}^{*})v_{i}}{rtp_{i}^{*}(1-p_{i}^{*})+p_{i}^{*}+h\left(\frac{1}{p_{i}^{*}}\right)}, \text{ for } i = 1, 2, \text{ and } \Upsilon(p_{1}^{*}) = 0, \text{ where}$$
$$\Upsilon(p_{1}) \stackrel{\text{def}}{=} \frac{\frac{w_{2}v_{2}^{t}}{w_{1}v_{1}^{tt}}p_{1}f\left[h\left(\frac{1}{1-p_{1}}\right),1\right]^{r}}{\left[rtp_{1}(1-p_{1})+1-p_{1}+h\left(\frac{1}{1-p_{1}}\right)\right]^{rt}} - \frac{(1-p_{1})f\left[h\left(\frac{1}{p_{1}}\right),1\right]^{r}}{\left[rtp_{1}(1-p_{1})+p_{1}+h\left(\frac{1}{p_{1}}\right)\right]^{rt}}.$$
$$\blacksquare \text{ The equilibrium is unique if } r\eta\left(\frac{1}{p_{i}}\right)\sigma\left(\frac{1}{p_{i}}\right) \leq 1.$$

J. Lagerlöf (U of Copenhagen)

Hybrid All-Pay and Winner-Pay Contests

June 5, 2018 19 / 24

・ロト ・ 母 ト ・ ヨ ト ・ ヨ ト … ヨ

- I assume n = 2 and I study three models:
 - The CSF is biased in favor of one contestant.
 - One contestant has a higher valuation than the other.
 - I also endogenize the degree of bias.
- Assumption 3. The CSF is given by

$$p_i(\mathbf{s}) = \frac{w_i s_i^r}{w_1 s_1^r + w_2 s_2^r}.$$

• The following three equations define equilibrium values of p_1^* , y_1^* , and y_2^* :

$$y_{i}^{*} = \frac{rtp_{i}^{*}(1-p_{i}^{*})v_{i}}{rtp_{i}^{*}(1-p_{i}^{*}) + p_{i}^{*} + h\left(\frac{1}{p_{i}^{*}}\right)}, \text{ for } i = 1, 2, \text{ and } \Upsilon(p_{1}^{*}) = 0, \text{ where}$$
$$\Upsilon(p_{1}) \stackrel{\text{def}}{=} \frac{\frac{w_{2}v_{1}^{\prime t}}{w_{1}v_{1}^{\prime t}}p_{1}f\left[h\left(\frac{1}{1-p_{1}}\right),1\right]^{\prime}}{\left[rtp_{1}(1-p_{1})+1-p_{1}+h\left(\frac{1}{1-p_{1}}\right)\right]^{\prime t}} - \frac{(1-p_{1})f\left[h\left(\frac{1}{p_{1}}\right),1\right]^{\prime}}{\left[rtp_{1}(1-p_{1})+p_{1}+h\left(\frac{1}{p_{1}}\right)\right]^{\prime t}}.$$

• The equilibrium is unique if $r\eta\left(\frac{1}{p_i}\right)\sigma\left(\frac{1}{p_i}\right) \leq 1.$

- A Biased decision process ($w_1 \neq w_2$ but $v_1 = v_2$)
 - Among the results:
 - (a) $p_1^* > p_2^* \Leftrightarrow y_1^* < y_2^* \Leftrightarrow C(s_1^*, p_1^*) > C(s_2^*, p_2^*).$ (b) Evaluated at symmetry $(w_1 = w_2): \frac{\partial p_1^*}{\partial w_1} > 0,$

$$\frac{\partial y_1^*}{\partial w_1} < 0, \quad \frac{\partial y_2^*}{\partial w_1} > 0, \quad \frac{\partial x_1^*}{\partial w_1} > 0 \Leftrightarrow \frac{\partial x_2^*}{\partial w_1} < 0 \Leftrightarrow \sigma(2) > \frac{2}{2 + rt}.$$

Different valuations ($v_1 \neq v_2$ but $w_1 = w_2$)

Among the results:

(a)
$$p_1^* > p_2^* \Leftrightarrow \frac{y_1^*}{v_1} < \frac{y_2^*}{v_2}.$$

(b) $v_1 - y_1^* > v_2 - y_2^* \Leftrightarrow C(s_1^*, p_1^*) > C(s_2^*, p_2^*)$

- A Biased decision process ($w_1 \neq w_2$ but $v_1 = v_2$)
 - Among the results:
 - (a) $p_1^* > p_2^* \Leftrightarrow y_1^* < y_2^* \Leftrightarrow C(s_1^*, p_1^*) > C(s_2^*, p_2^*).$ (b) Evaluated at symmetry $(w_1 = w_2): \frac{\partial p_1^*}{\partial w_1} > 0,$

$$\frac{\partial y_1^*}{\partial w_1} < 0, \quad \frac{\partial y_2^*}{\partial w_1} > 0, \quad \frac{\partial x_1^*}{\partial w_1} > 0 \Leftrightarrow \frac{\partial x_2^*}{\partial w_1} < 0 \Leftrightarrow \sigma(2) > \frac{2}{2 + rt}.$$

Different valuations ($v_1 \neq v_2$ but $w_1 = w_2$)

Among the results:

(a)
$$p_1^* > p_2^* \Leftrightarrow \frac{y_1^*}{v_1} < \frac{y_2^*}{v_2}$$
.
(b) $v_1 - y_1^* > v_2 - y_2^* \Leftrightarrow C(s_1^*, p_1^*) > C(s_2^*, p_2^*)$.

ヘロト 人間 とくき とくき とうせい

An Endogenous Bias (w_1 chosen, but $v_1 \ge v_2$ and w_2 fixed)

- Timing of events in the game:
 - **1** A principal chooses w_1 to maximize $R^H = C(s_1^*, p_1^*) + C(s_2^*, p_2^*)$.
 - 2 w_1 becomes common knowledge and the contestants interact as in the previous analysis.
- Assumption 3. The production function is of Cobb-Douglas form: $f(x_i, y_i) = x_i^{\alpha} y_i^{\beta}$, for $\alpha > 0$ and $\beta > 0$.

Results: The equilibrium values of p_1 and w_1 satisfy:

If
$$v_1 = v_2$$
, then $\widehat{p}_1 = \frac{1}{2}$ and $\widehat{w}_1 = w_2$.

If $v_1 > v_2$, then $\hat{p}_1 > \frac{1}{2}$.

If $v_1 > v_2$, then $\widehat{w}_1 < w_2$ at least if $|v_1 - v_2|$ is very small or big. My intuition for results:

- Contestant 1 is more valuable as a contributor (as $v_1 > v_2$).
- Hence, she should be encouraged to use x₁, as all-pay investments are more conducive to large expenditures.
- This is achieved by making winner-pay inv. costly: $\hat{p}_1 > \frac{1}{2}$.
- To generate p̂₁ > 1/2, v₁ > v₂ is more than enough, so bias can be in favor of Contestant 2.
 - Might not be robus

June 5, 2018 21 / 24

イロト イポト イヨト イヨト

An Endogenous Bias (w_1 chosen, but $v_1 \ge v_2$ and w_2 fixed)

- Timing of events in the game:
 - **1** A principal chooses w_1 to maximize $R^H = C(s_1^*, p_1^*) + C(s_2^*, p_2^*)$.
 - 2 w₁ becomes common knowledge and the contestants interact as in the previous analysis.
- Assumption 3. The production function is of Cobb-Douglas form: $f(x_i, y_i) = x_i^{\alpha} y_i^{\beta}$, for $\alpha > 0$ and $\beta > 0$.
- Results: The equilibrium values of p_1 and w_1 satisfy:

If
$$v_1 = v_2$$
, then $\widehat{p}_1 = \frac{1}{2}$ and $\widehat{w}_1 = w_2$.


If
$$v_1 > v_2$$
, then $\hat{p}_1 > \frac{1}{2}$.

If $v_1 > v_2$, then $\widehat{w}_1 < w_2$ at least if $|v_1 - v_2|$ is very small or big. My intuition for results:

- Contestant 1 is more valuable as a contributor (as $v_1 > v_2$).
- Hence, she should be encouraged to use x₁, as all-pay investments are more conducive to large expenditures.
- This is achieved by making winner-pay inv. costly: $\hat{p}_1 > \frac{1}{2}$.
- To generate $\hat{p}_1 > \frac{1}{2}$, $v_1 > v_2$ is more than enough, so bias can be in favor of Contestant 2.
 - Might not be robust.

Numerical example ($t = r = v_2 = w_2 = 1$)

Plot of plot p̂₁ and ŵ₁ against v₁ for three different values of α: 0.9 (the blue, dotted curve), 0.5 (the green, dashed curve), and 0.1 (the red, solid curve).



Main results and contributions: (1/1)

I The analytical approach (borrowing from producer theory):

 $\blacksquare \rightarrow$ Generality, tractability, and an existence condition.

A larger n leads to substitution away from all-pay investments.
But only if the elasticity of substitution is large enough.

3 Total expenditures always lower in hybrid contest than in all-pay.

4 Total exp'tures can be decreasing in n (also shown by Melkoyan).

 Asym. contests (in terms of valuations and bias): Sharp predictions about relative size of investm's and of expenditures.

6 Endogenous bias: High-valuation contestant more likely to win but the bias is against her (the latter might not be robust).

June 5, 2018 23 / 24

Possible avenues for future work (1/1)

- **1** Sequential moves: first (x_1, y_1) , then (x_2, y_2) .
- 2 Applying the producer theory approach to other contest models with multiple influence channels.

3 Experimental testing.

- Relatively sharp predictions.
- But risk neutrality might be an issue?
- 4 Further work on asymmetric contests.
 - More than two contestants.
 - Can a contestant be hurt by a bias in favor of her?
 - Can a contestant benefit from an increase in rival's valuation?

5 Contest design in broader settings.